

SPECIMEN LABEL

AC FORMULA 90[™] Ready-to-Use RODENTICIDE

No Prebaiting Required Just Use As Directed

Kills Norway Rats & Roof Rats and House Mice.
Rodents may consume a lethal dose in one feeding with first dead rodents appearing four or five days after treatment begins.

ACTIVE INGREDIENTS: Chlorophacinone,

2-[(p-chlorophenyl) phenylacetyl)]

 1, 3-indandione,
 0.005%

 INERT INGREDIENTS:
 99.995%

 TOTAL:
 100.000%

EPA REG. NO. 56-70 EPA EST. NO. 56-OH-1

KEEP OUT OF REACH OF CHILDREN CAUTION

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

1.5 OZ. BAG

Manufactured By: J.T. Eaton & Co., Inc., 1393 East Highland Road, Twinsburg, OH 44087

AC FORMULA 90^{TM} is a trademark of J.T. EATON & CO., INC. J.T. EATON $\!\!\!^{\text{\tiny B}}$ is a registered trademark of J.T. EATON & CO., INC.

PRECAUTIONARY STATEMENTS: HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Keep away from humans, domestic animals and pets. Keep away from feed or foodstuffs. May be harmful or fatal if swallowed or absorbed through the skin because this material may reduce the clotting ability of blood and cause bleeding.

NOTE TO PHYSICIÁN: If ingested, administer Vitamin K1 intramuscularly or orally, as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Do not apply this product directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store only in original container, in a cool, dry place

inaccessible to children and pets.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty container. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned stay out of smoke.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help to prevent accidents:

- 1. Store product not in use in a location out of reach of children and pets.
- 2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
- 3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: For control of Norway Rats, Roof Rats and house mice in and around homes, industrial and agricultural buildings, and similar man-made structures. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS: Determine areas where rats and mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much food as possible. **APPLICATION DIRECTIONS:**

Rats: Apply 3 to 10 place packs per bait placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

Mice: Apply one place pack per bait placement. Space place packs at 8- to 12- foot intervals. Maintain an uninterrupted supply of fresh bait for at least 15 days.

Rats and Mice: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.